MISSISSIPPI MAIL SURVEY OF GAME HARVEST 1971 - 72

STUDY COMPLETION REPORT FEDERAL AID PROJECT NO, W-48-R STUDY VI

BY

BILL QUISENBERRY, JR.

AUGUST 29, 1972



GAME DIVISION

AN ANALYSIS OF MISSISSIPPI'S FIRST POST-SEASON GAME HARVEST MAIL SURVEY (1970-71)

Ву

WILLIAM YOUNG QUISENBERRY III

APPROVED:

Professor of Wildlife and Fisheries and Head of the Department (Chairman of Supervisory Committee)

Assistant Professor of Wildlife and Fisheries (Member of Committee)

Associate Professor of

Bctany (Minor Professor) Dean of the School of Forest Resources

Assistant Professor of Wildlife and Fisheries (Member of Committee)

Dean of the Graduate School

ACKNOWLEDGMENTS -

I wish to acknowledge the Mississippi Game and Fish Commission for making funds available for this project.

The advice, suggestions, and assistance in other forms of Mr. John L. Given, Jr., Head of the Division of Data Processing, Mississippi Game and Fish Commission, is deeply appreciated. I am also indebted to Mrs. Ann Faries, Mrs. Barbara Bowering, Mrs. Gloria Hales, and Mrs. Dianne Guinn of the Data Processing Division, who assisted in the programming, key punching, and verification of the survey data.

Special thanks are extended to Mr. Harold W. Rimmer, Head of the Accounting Division, Mississippi Game and Fish Commission, and his staff for assistance in the preparation of the questionnaires for mailing.

A note of appreciation is expressed to Dr. Don W.

Hayne, Institute of Statistics, North Carolina State

University, for his invaluable guidance and constructive

criticism. The technical assistance and advice of Mr.

Robert E. Mason and Mr. David W. Turner, also associated

with the Institute of Statistics, is gratefully acknowledged.

Special thanks are due my graduate committee for their helpful suggestions in reviewing the manuscript. The committee consisted of Dr. Dale H. Arner, Head of the Department of Wildlife and Fisheries and Director of the Thesis; Dr. Wendell J. Lorio, Department of Wildlife and

Fisheries; Dr. J. Ray Watson, Department of Botany; and.
Mr. Bobby R. Tramel of the United States Department of
Interior, Bureau of Sport Fisheries and Wildlife.

W. Y. Q.

TABLE OF CONTENTS

	age
ACKNOWLEDGMENTS	iii
LIST OF FIGURES	vi
LIST OF TABLES	vii
INTRODUCTION	1
METHODS AND PROCEDURES	6
RESULTS AND DISCUSSION	15
SUMMARY	20
ABSTRACT	23
APPENDIX	25
LITERATURE CITED	49

LIST OF. FIGURES -

Figu	ure	-	<i>:</i> .										· P	'age
1.	Letter of	Transn	nittal	- F	irst l	Maili	ng	٠		٠	٠		19	10
2.	Letter of Third Mai						. •		d	٠	٠	٠	•	11
3.	State of D Questionn				-71 H									12
4.	Distributi Communiti Are Indica	es Repr	esente	dii	n the	Sam	ple	, its	w	. 6		٠	•	13
5	Mississin	of Game	Manag	eme	nt Di	stric	ets						, ,	14

LIST OF TABLES

Tab	<u>Page</u>	
1.	State-Wide Game Kill, 1950-51 Season 26	
2.	1970-71 Survey Mailing and Response Data 27	
3.	Summary 1970-71 Mississippi Post-Season Game Harvest Mail Survey	
4.	Summary Dove Harvest Mail Survey 1970-71 29	
5.	Summary Bobwhite Quail Harvest Mail Survey 1970-71	
6.	Summary Rabbit Harvest Mail Survey 1970-71 31	
7.	Summary Deer (Archery Season) Harvest Mail Survey 1970-71	
8.	Summary Deer (Gun Season) Harvest Mail Survey 1970-71	
9.	Summary Deer by Either Bow or Gun (Hunters not Additive) Harvest Mail Survey 1970-71	
10.	Summary Turkey Harvest Mail Survey 1970-71 35	
11.	Summary Squirrel Harvest Mail Survey 1970-71	
12.	Summary 'Duck Harvest Mail Survey 1970-71	
13.	Summary Woodcock Harvest Mail Survey 1970-71	
14.	County Summary' of Hunting Pressure	
15.	District Summary of Hunting Pressure 41	
16.	Licensees Who Have Hunted or Trapped Other Species in Mississippi	
17.	Licensees Who Have Hunted or Trapped Other Species in Mississippi - Expanded Summary	

sorted into sub-sample sequence and returned to the computer system with the 5081 cards and the effective sample size (number of questionnaires mailed minus the number returned by the post office as non-deliverable) for each sub-sample was calculated.

Licensees were asked to furnish information concerning number of days afield, and total number harvested for the following species: mourning dove, bobwhite quail, rabbit, squirrel, duck, woodcock, deer, and turkey. In addition, licensees were requested to indicate which of the following species they have hunted or trapped in Mississippi: fox, crow, raccoon, opossum, beaver, bobcat, and geese. An attempt was made to determine general hunting pressure by county and game management district (Figure 5). Recipients were asked if they would purchase a user permit, advocate the establishment of more public hunting areas, and whether or not hunting activity occurred on a state or federal hunting area.

Mississippi utilized the same general game harvest mail survey procedures as do Maryland (Hodil, personal communication 1968, 1971), Tennessee (Legler, personal communication 1971), Alabama (Kelly, 1967, 1968), North Carolina (Barick and Critcher, 1955), Missouri (Sampson, 1965, 1968, 1969), Michigan (Eberhardt, 1961), New Jersey (Smith, 1968; Wright, 1964), New York (Maguire, 1962), and Kentucky (Durell, 1967).

Game and Fish Commission

COMMISSIONERS

P. O. BOX 451 • PHONE 354-7333 • JACKSON, MISSISSIPPI 39205

COOL. JR. CHAIRMAN LUCEDALE

JOHN BELL-WILLIAMS GOVERNOR

BILLY JOE CROSS

CHARLES L. SULLIVAN LIEUTENANT GOVERNOR

COMMISSIONERS

R COVINGTON MERIDIAN

TUPELO

SAMIL

GULFPORT

BRENT NICKLE

OXFORD

TOM RIDDELL, JR CANTON

L. H SANDERS CARTHAGE

WATER VALLEY

MARSHALL D. TREPPENDAHL

GLEVELAND TOM W. CLEVELAND JACKSON

VICE CHAIRMAN

WOODVILLE

Fellow Sportsman:

have been selected as a representative hunter to furnish us information concerning the 1970-71 hunting season

Your reply on the enclosed form is very important. For accurate results, this information is needed from everyone receiving a question-naire.' If you did not hunt during 1970-71, please answer question one and return the questionnaire. DO NOT pass the questionnaire on to a friend who did hunt.

The information supplied by you and other selected hunters will assist the Mississippi Game and Fish Commission in managing our game resources and in wisely administering license fee funds. It is neces sary that we make periodic checks in order to measure the effectiveness of our regulations and management practices.

Only a limited number of hunters can be contacted so a good response is needed for reliability.

As one of the hunters selected, you are requested to fill in the enclosed questionnaire as accurately as possible and return it in the self-addressed envelope provided.

A prompt reply will be sincerely appreciated.

Yours for better conservation,

Billy Joe Cross Executive Director

BJC:nj

Figure 1. Letter of Transmittal - First Mailing

Game and Fish, Commission

STATE OF MISSISSIPPI

COMMISSIONERS

P. O. BOX 451 . PHONE 354-7333 . JACKSON, MISSISSIPPI 39205

R. G. COOK, JR CHAIRMAN LUCEDALE

JOHN BELL WILLIAMS

MARSHALL D TREPPENDAHL VICE CHAIRMAN WOODVILLE

MERSHEL & HOWELL WATER VALLEY

J. E WOLFE

TOM W. CLEVELAND



CHARLES L. SULLIVAN LIEUTENANT GOVERNOR COMMISSIONERS

BILL" R. COVINGTON MERIDIAN

RIRBY P. FAUCETTE

SAM V. MORSE GULFPORT

DDENT NICKEE

BRENT NICKLE DXFORD TOM RIDDELL, JR.

L. H. SANDERS

CANTON

Fellow Sportsman:

Recently we mailed you a Hunter Questionnaire and requested that you fill out and return the completed form. In case you have misplaced the original form, or haven't had an opportunity to canplete it as yet, we are enclosing a duplicate and request that you take a few minutes to complete and return it in the enclosed stamped envelope.

We need your response to assure a high percentage of returns which are necessary because the number of hunters we are contacting is relatively small.

The **information** requested **from** you is for the purpose of **managing** our game resources. Please fill out the farm and return it even if you did not hunt, or **were** not very successful in your efforts.

Your cooperation will assist us in better directing the management of Mississippi's game resources.

Thank you.

Yours for better conservation,

Billy Joe Cross Executive Director

Enclosures

Figure 2. Letter of Transmittal - Second and Third Mailings

STATE OF MISSISSIPPI 1970-71 HUNTER QUESTIONNAIRE

										VEC	NO
FOX	CROW	RACCOON	OPPOSSUM	BEAVER	ВО	BCAT	GEESE	OTHE	R O	THER	OTHER
TRUCT	IIONS: PLE				T YOU	HAVE	HUNTED OR		M SS	SS PP	I.
		col	JNTÝ	DAYS					COUNTY		DAYS
		COL	JNTY	_DAYS					COUNTY	*********	DAYS
		COL		DAYS					COUNTY		_DAYS
) ROCI	NUM	ABER OF DAY	'S HUNTED	IN EACH C	OUNTY	119 1111	100 11	OHTED.			
701107		ASE LIST T	THE MISSISS								
DID A		TURKEY LAST	SPRING?				RIL 26, 1 DAYS? —				,
	(B), GU1	SEASON? Y	ÆS — NO —	<u> </u>			DAYS?	TOTAL	KILL FOR	SEASO	N
		CHERY SEASO									
010 Y											
DID Y	YOU HUNT \	WOODCOCK?	/ES NO -		ON HOW	V MANY	DAYS?	_ TOTAL	KILL FOR	SEASON	I ——
DID Y	YOU HUNT I	DUCKS? YFS_	NO		ON HO	W MANY	DAYS?	_ TOTAL	KILL FOR	SEASON	
DID Y	OU HUNT	QUIRREL'S?	YES NO		ON HO	MANY	DAYS* —	- TOTAL	KILL FOR	SEASON	
DID Y	OU HUNT !	RABBITS? YE	5 NO	-	ON HO	V MANY	DAYS?	_ TOTAL	KILL FOR	SEASON	
DID Y	YOU HUNT (30BWHITE QU	JATL? YES—	— NO	OH HO	W MANY	DAYS" —	— TOTAL	KILL FOR	SEASON	
010	YOU HUNI !	MOURNING DO	OVES? YFS	NO	ON HO	W MANY	DATE:	_ IOTAL	KILL FOR	SEASON	
010	AND DESIGNATION	ANY GAME DU	INTING THE	070 71 H	UNITEN	C CEAS	ON THE MIC	CICCIPPI	2 YF5	NO	
						AND RE	TURN THIS	FORM.			
							PLEASE A			,	
£51	POINT			MS 397			UNT IT AS				_
1 2	BUX 70						HAN ONE K				
	TEICAT	ION NO.	3-s JOHN R		1	HIGHT	HAVE REEN	WITH YO	U. IF YO	U HUNTI	E,D
						DO NOT	REPORT I	HE KILL	OF ANYON	= Maritis	

Figure 3. State of Mississippi 1970-71 Hunter Questionnaire




Figure 4. Distribution of the Survey Sample - Communities Represented in the Sample Are Indicated by Dots




Figure 5. Mississippi Game Management Districts

RESULTS AND DISCUSSION'

This analysis is based on the total response from three mailings at two-week intervals each from which 4,142 questionnaires (91.2 percent) were presumed delivered by the post office (402 or 8.8 percent were returned as undeliverable). One thousand eight 'hundred ninety-one (42 percent) of the selected licensees responded to the first mailing; 922 (20 percent), the second mailing; 394 (09 percent), the third mailing. Nine hundred thirty-five questionnaires (22.6 percent) had not been returned when the survey was terminated.

Using delivered 'questionnaires as a base, the 3,207' returns represented a response of 77.42 percent and the usable returns (3,205), a response of 77.38 percent. The usable returns constituted a 1.33 percent sample of Type I and Type II licenses purchased during the 1970-71 season. Table 2 is a summary of the survey mailfng and response data.

Of the 3,207 licensees who' returned questionnaires,

2,696 (84.12 percent) sought game during the season. Five
hundred nine respondents (15.75 percent) did not 'hunt after
purchasing a license. State-wide, an estimated 203,260
(84.52 percent) of Mississippi's 240,479 resident licensed
hunters participated in some form of game hunting recreation.

A summary of the survey results is presented in Table 3. Data reveal squirrel to be the most popular game species hunted (62 percent), followed by 'deer - gun season (50 per-'cent), rabbit (38 percent), mourning dove (35 percent), bobwhite quail (32 percent), duck (15 percent), turkey. (10 percent), deer - archery season (05 percent), and woodcock '(03 percent). Respondents harvested 30 Squirrel. 30,052 mourning dove, 29,834 bobwhite quail, 13,867. rabbit, 3,649 duck, 727 woodcock, 527 deer (gun season), 24 deer (archery season), and 94 turkey. Mourning do 744 hunters were the most successful (96 percent), followed by squirrel (94 percent), bobwhite quail (92 percent), rabbit and woodcock (91 percent), duck (82 percent), turkey (24 percent), deer gun season (23 percent), and deer archery season (14 percent).

Information pertaining to each species consists of number hunters, total days afield, total harvest, average daily bag, average season bag, and average days afield for each of the th'ree mailings (Tables 4-13). A summary, of mourning dove data is found in Table 4; bobwhite quail,' Table 5; rabbit, Table 6; deer = archery season, Table 7; deer = gun season, Table 8; deer = hunters not additive,. Table 9; turkey, Table 10; squirrel, Table 11; duck, Table 12; and woodcock, Table 13.

Hunting pressure, based on 2,633 usable responses (82.15 percent), was determined by county (Table 14) and

game aanagement district (Table 15). Results indicated that respondents spent 54,124 days in the field, an average of 4.8 days per hunter. Total hunter response (4,860) would suggest that many licensees hunted in more than one county during theseason. Jones, Jackson, Bolivar, Lafa-yette, Pike, Kemper, Yazoo, Lauderdale, Monroe, Scott, and. Warren counties, respectively, absorbed the greatest hunting pressure. Most hunting activity was centered in District VI (19.76 percent), while District III received the least (14.58 percent).

Data from this segment of the Survey were not statistically analyzed, and a limited number of usable responses were tabulated. However, a general state-wide pattern of hunting pressure was established and useful, reasonably accurate information can be derived from this data.

Table 16 is a summary of licensees who have hunted or trapped other species in Mississippi, based on 3,205 usable responses. Raccoon was the most popular sought after game as indicated by the Sampled hunters (15.28 percent), followed by fox (8.42 percent), crow (8.11 percent), opossum (7.50 percent), bobcat (4.41 percent), beaver (2.28 percent, and geese (1.39 percent). Table 17 points out the expanded estimate of licensees who have 'hunted or trapped other species in Mississippi. Confidence limits at the 95 percent probability level for this expanded estimate are listed in Table 18.

Two thousand four hundred seventy respondents (77.03 percent) did not hunt on state or federal hunting areas during the season. Hunter sentiment (73.60 percent) would suggest that Mississippi should establish more public hunting areas. One thousand seven hundred seventy-six sampled hunters (55.41 percent) would purchase a user permit, in addition to hunting license, to make available additional facilities or services on public hunting areas.

In regard to the user permit, many hunters expressed confusion and indecision. Comments such as "don't understand the question," "If the price is right," "license costs too much already," "don't know what a user permit is," and "too little information available to make a decision," were not uncommon. In retrospect, this was a poorly constructed question, and the accuracy of this particular data is questionable.

State-wide, an estimated 176,735 licensees (73.49 percent) did not hunt on public land; 63,744 (26.51 percent), did. An estimated 214,499 individuals (89.20 percent) favor the establishment of more public hunting areas in Mississippi; 259,699 (66.41 percent) would purchase a user permit in addition to a hunting license.

An expanded estimate including number of hunters, percentage of licensees, total days afield, total harvest, average daily bag, average season bag, and average days afield for eight game species is presented in Table 19.

State-wide, these eight species provided Mississippi hunters

an estimated 4,034,735 days afield with an estimated harvest of 8,017,191 individual items of game. Standard error as applied to estimated number of hunters is tabulated in Table 20; days afield, Table 21; harvest, Table 22. Statewide confidencelimits at the 95 percent probability level for estimated number of hunters, total days afield, and total harvest, are pointed out in Table 23 and indicate a reasonable degree of high reliability (Drapala, personal interview 1971).

This initial project is a success and should be continued annually for a period of at least five years in order that Mississippi's game resources might be more effectively managed.

SUMMARY

- 1. Two thousand six hundred thirty-three respondents spent 54,124 days afield, an average of 4.8 days per hunter.
- 2. Jones, Jackson, Bolivar, Lafayette, Pike, Kemper, Yazoo, Lauderdale, Monroe, Scott, and Warren counties, respectively, absorbed the heaviest hunting pressure. Most hunting activity was centered in District VI (19.76 percent), while District III received the least (14.68 percent).
- 3. Squirrel was the most popular game sought by the respondents, (62 percent), followed by deer = gun season (50 percent), rabbit (38 percent), mourning dove (35 percent), bobwhite quail (32 percent), duck (15 percent); turkey (10 percent), deer = archery season (05 percent), and woodcock (03 percent).
- 4. Respondents harvested 30,344 squirrel, 30,052 mourning dove, 29,835 bobwhite quail, 13,867 rabbit, 3,649 duck., 727 woodcock, 527 deer (gun season), 24 deer [archery season), and 94 turkey.,
- 5. Mourning dove hunters were the most successful (96 percent), followed by squirrel (94 percent), bobwhite quail (92 percent), rabbit and woodcock (91 percent), duck (82 percent), turkey (24 percent), deer gun, season (23 percent), and deer archery season (14 percent).

Previously, only two game surveys have been carried out. Leopold (1929) completed the first game survey of Mississippi. No estimated game harvest data were compiled in this publication. Thompson and Redmond (1951) were responsible for an analysis of game harvest during the 1950-51 hunting season (Table 1). Approximately 40 percent. of 10,000 licensed hunters received a questionnaire (Hunters' Scorecard). Information supplied by the respondents was not statistically analyzed.

METHODS AND PROCEDURES

Mississippi Game and Fish Commission personnel conducted the state's first post-seasongame harvest mail survey during 1971. The Institute of Statistics, North Carolina State University, was contracted to complete a statistical analysis of the detail data.

Licensee names and addresses were selected at random from the current files of Type I (combination hunting and. fishing) and Type IS (hunting only) license stubs. License numbers with the following terminal digits provided by Dr. Don W. Hayne, Institute of Statistics, North Carolina State University, were selected: 017, 018, 019, 020, 409, 410, 411, 412, 593, 594, 595, 596, 797, 798, 799, 800, 909, 910, 911, 912 (Hayne, personal communication 1970). Of 240,479 Type I and Type II licenses purchased during the 1970-71 season, 4,544 individuals (1.89 percent) were selected to participate in the survey. Data supplied by 3,205 respondents (1.33 percent) were included in the analysis.

The initial mailing of questionnaires was made March 15, 1971, after the close of bobwhite quai'l season. Two regular follow-up mailings to non-respondents were made on March 31, 1971 (2,359), and April 16, 1971 (1,491). The survey was terminated May 4, 1971.

Selected hunters received a letter of transmittal (Figures 1 and 2), questionnaire form (Figure 3), and a self-addressed, postage-paid business reply envelope in a printed bulk mail permit window envelope. First class mail was used in all mailings.

Although no deliberate attempt to equalize geographic distribution was made, licensees residing in each of Mississippi's 82 counties were represented in the randomized sample (Figure 4). Communities represented in the sample are indicated by dots.

An IBM 360-20 computer system was utilized during the study. The name and address of each licensee was punched into a standard general purpose card. An identification number assigned to each licensee was created at the same time a master name and address tape file was written onto magnetic tape. Identification numbers were punched into the alphabetically arranged cards and printed on the questionnaire forms as a means of separating respondents from nan-respondents and delivered from undelivered forms for subsequent mailings. Separation was accomplished by punching identification numbers of respondents on finder cards and machine-matching with the address cards to remove respondents from the address card deck (Sampson, 1965).

The continuous printed questionnaires were designed as three-part color-coded forms (first mailing, white; second mailing, green; third mailing, pink), and all were machine printed at one time.

Responses were edited to determine credibility and were coded to facilitate key punching and verification. Forms containing any valid data were considered usable by the writer. Questionnaires returned by a family member indicating that the selected licensee had moved, was in military service, had deceased, etc., were considered undelivered rather than unusable.

Usable responses were delivered to the key punch operator who recorded only the identification number in general purpose cards. The cards were sorted into ascending sequence and matched against the master name and address file to be updated and to determine which questionnaires were to be included as valid input for detail data. Valid questionnaires were returned to the key punch operator who recorded the detail data in general purpose cards. At the close of each mailing the identification numbers as recorded on the updated master file were sorted to determine which questionnaires should be included in the next mailing. Identification numbers of undelivered questionnaires were key punched so that the master file could be updated as they were returned. Deletion cards for undelivered questionnaires were held until the close of t e third mailing.

At the close of the third mailing, the original master file was returned to the computer system and the number of respondents in each of five sub-samples was calculated and punched into 5081 cards. Undelivered deletion cards were

LIST OF TABLES (Continued)

Tabl				Pa	age
18.	Confidence Limits at 95 Percent Probability Level - Expanded Estimate of Licensees Who Have Hunted or Trapped Other Species in Mississippi (Variation from the Mean)	•		•	44
19.	Expanded Summary 1970-71 Mississippi Post-Season Game Harvest Mail Survey	•		•	45
20.	Mississippi Post-Season Game Harvest Mail Survey 1970-71 - Expanded Estimate of Total Hunters	•			46
21.	Mississippi Post-Season Game Harvest Mail Survey 1970-71 - Expanded Estimate of Total Days Afield	•			47
22.	Mississippi Post-Season Game Harvest Mail Survey 1970-71 - Expanded Estimate of Total Harvest	•	•		48
23.	Confidence Limits at the 95 Percent Probability Level - Expanded Estimate of Total Hunters, Total Days Afield, Total Harvest (Variation from the Mean)		•		49

Wildlife regulatory agencie have for some time been approaching a system of adjustable regulations which vary with the status of the species, often on an annual basis. This system requires periodic estimates of relative population levels and harvests over extensive areas. These estimates can be obtained only through sampling surveys. Within the last several years considerable attention has been focused on such surveys throughout the Southeastern United States. As a result, call'counts, road counts, bag counts, and other sampling techniques were developed (Overton, 1955).

Although specific objectives and techniques for ob-, taining the desired information have varied, the primary purpose has been to obtain basic data upon which improved wildlife management practices can be based. Barick and. Critcher (1955) suggest five functions of the game survey that supply these basic data:

- (1) The determination of the relative immortance of individual game species as reflected by the amount of hunter-effort and the annual kill.
- (2) The determination of hunter-success which is, to some degree, an index to availability and relative abundance.
- (3) The correlation of annual trends in kills, hunter-success, and hunter-effort with annual variations in natural factors, such as weather and food supplies.

(4) In a similar manner, an evaluation and correlation of the effects of natural catastrophes, such as floods, fires, and hurricanes.

(5) Correlations between kills and annual changes in hunting seasons and bag limits.

Fluctuations in harvest from one year to the next should be the primary concern of the game harvest analysis. These fluctuations may reflect a change in the population level or they may be largely a function of other factors, such as weather, seasonal availability, hunting pressure influenced by some outside force, or the survey technique itself. Therefore, the nature and cause of these fluctuations must be carefully analyzed (Barick and Critcher, 1955).

Mail surveys have assumed an important role in the present administration of game and fish, primarily because they are the most inexpensive method of obtaining information that may be evaluated with existing statistical techniques (Overton, 1955). These surveys, however, may lead to bias.

Several important sources of bias are present. O these, sampling error, response error, and non-response bias are the most common. It is felt that such biases 'are more serious and more difficult to Isolate and correct in the field of game harvest surveys than in some other types of surveys (Hiltunen, 1952; Barick and Critcher, 1955; Martinson and Whitesell, 1964; Hayne, personal communication 1968).

Sampling error results when the sample of contacts is not representative of the subject of interest and is probably minimal when proper sampling 'techniques are used (Chapman., et al., 1959). Response errors in mail questionnaire surveys of hunters have been recognized as one of the major problems in that method of obtaining hunting statistics (Hiltunen, 1952; Atwood, 1956; Cronan, 1960; Hayne, 1964).

Hiltunen (1952), Overton (1955), Martinson and Whitesell (1964), and Hayne (1964) among others, have found that hunters tend to report more hunting activity and better success than they actually experienced with the response errors thus incurred resulting in a positive bias affecting the data collected.

"Prestige Bias" and "Memory Bias" are the two most apparent response errors (Overton, 1955; Atwood, 1956; Cronan, 1960). Prestige bias is the term applied to the hesitancy to admit to poor success. Memory bias presumably occurs when a respondent guesses his seasonal kill. Few people remember exactly how much game they killed, and the greater the time lapse between the event and the estimate, the greater the bias is likely to be. Too, there is a natural tendency to suppress information on game taken illegally.

Non-response bias is due to the tendency for the more successful persons to answer the questionnaire more readily than the less successful ones. This phenomenon has been

- 10 Sp

4

Be.

Iţ.

12.30

45. - 1

12

36

suspected by many biologists and demonstrated by Hayne (1964) and Martinson and Whitesell (1964) among others.

Confidence intervals are an objective measure of reliability employed to ascertain the degree of confidence with respect to the accuracy of game harvest estimates. Narrowing the spread of confidence intervals would enable one to place more reliance on the data derived. Much greater faith could be placed on an estimate if confidence intervals could be held to not more than 10 percent of the computed kill (Barick and Critcher, 1955; Drapala, personal interview 1971). Further research in testing and evaluating confidence intervals as they apply to this type of survey is needed.

It may be stated that "although the game harvest mail survey shows much promise as a valuable game management tool, much work still needs to be done in the way of refining the technique. This refinement should be aimed at evaluating and compensating for biases and improving the confidence intervals, which, in turn, may be expected to produce more accurate data on kill fluctuations and population levels" (Barick and Critcher, 1955)

The purpose of the 1970-71 game harvest mail survey is to compile the first statistically reliable estimate of Mississippi's annual game harvest. It is also the first survey of its kind conducted in the state.

- 6. In regard to other species hunted or trapped in Mississippi, raccoon was sought by 15.28 percent of the sampled hunters, followed by fox (8.42 percent), crow (8.11 percent), opossum (7.50 percent), bobcat (4.41 percent), beaver (2.28 percent), and geese (1.39 percent).
- 7. State-wide, an estimated 203,260 (84.52 percent) of Mississippi's 240,479 resident licensed hunters participated in some form of game hunting recreation.
- 8. An expanded state-wide estimate of total number hunters, total days afield, and total harvest respectively for each kind of game follows: mourning dove 81,904 357,394 2,213,530; bobwhite quail 75,411 639,427 2,173,500; rabbit 91,442 702,804 1,059,120; deer (archery season) 10,296 62,543 1,608; deer (gun season) 118,935 853,984 35,886; deer by either bow or gun (hunters not additive) 119,939 911,803 37,353; turkey 22,548 103,165 6,355; squirrel 151,046 1,134,180 2,298,120; duck 33,217 159,080 254,236; woodcock 6,844 26,839 54,862.
- 9. State-wide, these eight species provided Mississippi hunters an estimated 4,034,735 days afield with an estimated harvest of 8,017,191 individual items of game.
- 10. Confidence intervals at the 95 percent probability level were computed state-wide for number of licensees

hunting each species, total days afield, and total harvest and indicate a reasonable degree of high reliability.

11. An estimated 89.20 percent of Mississippi hunters favor the establishment of more public hunting areas in the state; most did not hunt on state or federally controlled land (73.49 percent); 66.41 percent would purchase a user permit, in addition to a hunting license, to make available additional facilities or services on public hunting areas.

ABSTRACT

William Young Quisenberry, III, Master of Science, 1971

Major: Wildlife Ecology, .Department of Wildlife and

Fisheries

Title of Thesis: An Analysis of Mississippi's First

Post-Season Game Harvest Mail Survey

125

譹

A PLAN

100

报:

-19

A 77

(1970-71)

Directed by: Dr. Dale H. Arner

Pages in Thesis: 51 Words in Abstract: 287

ABSTRACT

Mississippi's first post-season game harvest mail survey was conducted during 1971. Names and addresses of 4,544 licensees were selected at random from the current file of 240,479 resident hunting- and combination hunting and fishing license stubs. An initial mailing with two follow-up mailings were used and consisted of a questionnaire, letter of transmittal, and business reply envelope. Data were programmed, compiled, key punched, and verified by personnel of the Data Processing Department, Mississippi Game and Fish Commission. Detail cards were mailed to the Institute of Statistics, North Carolina State University, where a statistical analysis was conducted.

General hunting pressure (number hunters and percent of hunting activity) by county and game management district was determined. State-wide, an estimated 203,260 licensees participated in some form of game hunting activity'.

An expanded state-wide estimate of total number hunters, total days afield, and total harvest, respectively, for eight game species follows: mourning dove 81,904 - 357,394 - 2,213,530; bobwhite quail 75,411 - 639,427 - 2,173,500; rabbit 91,442 - 702,804 - 1,059,120; deer (archery season) 10,296 - 62,543 - 1,608; deer (gun season) 118,935 - 853,984 - 35,886; deer by either bow or gun (hunters not additive) 119,939 - 911,803 - 37,353; turkey 22,548 - 103,165 - 6,355; squirrel 151,046 - 1,134,180 - 2,298,120; duck 33,217 - 159,080 - 254,236; woodcock 6,844 - 26,839 - 54,862.

Confidence intervals at 95 percent probability level were computed state-wide for number of licensees hunting each species, total days afield, and total harvest, and they indicate a reasonable degree of high reliability.

Mississippi hunters favor the establishment of more public hunting areas in the state (89.20 percent); most did not hunt on state or federally controlled land (73.49 percent); 66.41 percent would purchase a user permit, in addition to a hunting licenase, to make available additional facilities or services on public hunting areas.




Table 1. State-Wide Game Kill, 1950-51 Season

Species	Percentage Licensees Hunting	Number Hunters	Average No. Times Hunted	Average Kill Per Hunt	Average Kill Per Season	Estimated Harvest
Deer*	6.00	11,283			.12	1,370
Turkey	1.35	2,557	3.40	.028	.097	249
Quail	47'. 12	89,056	7.15	3.32	23.80	2,119,532
Squirrel	79.83	150,878	7.23	2.21	16.03	2,418,574
Rabbit	47.15	89,113	6.44	1.57	10.14	900,360
Raccoon	16.20	30,693	6.15	1.10	6.80	208,712
Opossum	23.06	43,583	5.16	1.71	8.86	386,145
Dove	21.58	40,786	3.25	3.9 3	12.82	,522,876
Duck	8.97	16,226	4.75	2.35	11.23	,182,217

^{*}Estimates based on Wardens reports on deer kill and deer camp permit holders reports.

Table 2. 1970-71 Survey Mailing and Response Data

First Mailing	Number	Total Percent
Total Mailed Undelivered Delivered Returned Usable Not Usable Not Returned	4,544 331 4,213 1,854 1,854 0 2,359	100.0% 7.2% 92.8% 44.0% 100.0%
Second Mailing	Number	Total Percent
Total Mailed Undelivered Delivered Returned Usable Not Usable Not Returned	2,359 50 2,309 922 922 0 1,387	100.0% 2.2% 97.8% 39.9% 100.0%
Third Mailing	Number	Total Percent
Total Mailed Undelivered Delivered Returned Usable Not Usable Not Returned	1,491 21 1,116 394 392 2 722	100.0% 1.4% 98.6% 35.0% 99.9% 0.5% 64.7%
Total	Number	Total Percent
Total Mailed Undelivered Delivered Returned Usable Not Usable Not Returned	4,544 402 4,142 3,207 3,205 02 935	100.0% 8.8% 91.2% 77.4% 77.3% 0.062% 22.6%

Table 3. Summary 1970-71 Mississippi Post-Season Game Harvest Mail Survey

Total Number Usable Replies: 3,205

(Number Who Actually Hunted: 2,697)

Species	Total Hunters	Percent Successful Hunters	Total Days Afield	Total Harvest	verage Daily Bag	Average Season Bag	Average Days Afield	Percentage of Licensees Who Hunted
Mourning Dove	1,115	96%	5,056	30,052	5.95	26.95	4.54	35%
Quail	1,048	92%	8,437	29,834	3.54	28.47	8.05	32%
Rabbit	1,203	91%	9,190	13,867	1.51	11.53	7.64	38%
Deer- Archery	150	14%	953	24	0.025	0.16	6.35	05%
Deer-Gun	1,607	23%	11,525	527	0.046	0.33	7.17	50%
Turkey	315	24%	1,455	94	0.065	0.30	4.62	10%
Squirrel	1,998	94%	14,808	30,344.	2.05	15.19	7.41	62%
Duck	480	82%	2,283	3,649	1.60	7.60	4.76	15%
Woodcock	100	91%	407	727	1.79	7.27	4.07	0 3%
		* .						

Summary Dove Harvest Mail Survey 1970-71 Based on 3,205 Usable Responses Table 4.

Mailing	Number Hunters	Total Days Afield	Total Harvest	Average Daily Bag	Average Season Bag	Average Days Afield	
First	685	3,310	19,456	5.88	28.40	4.83	
Second	309	1,216	7,286	5.99	23.58	3.94	
Third	121	530	3,310	6.25	27.36	4.38	
Total	1,115	5,056	30,052	5.94	26.95	4.54	

Table 5. Summary Bobwhite Quail Harvest Mail Survey 1970-71

Based on 3,205 Usable Responses

Number Hunters	Total Days Afield	Total Harvest	Average Daily Bag	Average Season Bag	Average Days Afield
630	5,487	18,613	3.39	29.57	8.75
302	1,916	7,582	3.96	25.11	6.34
116	1,034	3,639	3.52	31.37	8.91
1,048.	8,437	29,834	3.54	28.47	8.05
	Hunters 630 302 116	Number Days Afield 630 5,487 302 1,916 116 1,034	Number Hunters Days Afield Total Harvest 630 5,487 18,613 302 1,916 7,582 116 1,034 3,639	Number Hunters Days Afield Total Harvest Daily Bag 630 5,487 18,613 3.39 302 1,916 7,582 3.96 116 1,034 3,639 3.52	Number Hunters Days Afield Total Harvest Daily Bag Season Bag 630 5,487 18,613 3.39 29.57 302 1,916 7,582 3.96 25.11 116 1,034 3,639 3.52 31.37

	Table 6.	Summary Rabbit Based on	ubbit Harvest Mailed on 3,205 Usable	Mail	Survey 1970-71 Responses	
				- 1		
Mailing	Number Hunters	Total Days Afield	Total Harvest	Average Daily Bag	Average Season Bag	Average Days Afield
1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	069	5,351	8,015	1.50	11.67	7.76
Second	357	2,454	3,955	1.61	11.08	6.87
Third	156	1,385	1,897	1.37	12.16	88
Total	1,203	9,190	13,867	1.51	.11.53	7.64

Summary Deer (Archery Season) Harvest Mail Survey 1970-71 Table 7.

Based on 3,205 Usable Responses

					, . 1:1
Average Days Afield	9.0	5.42	6.19	6.35	
Average Season Bag	0.17	£0 °0	0.19	0.16	
Average Daily Bag	0.025	0.022	0.030	0.025	
Total	17	4		24	
Total Days Afield	675	179	6	953	
Number Hunters	101	Ж	16	150	
Mailing	First	Second	Third	Total	

Table 8. Summary Deer (Gun Season) Harvest Mail Survey 1970-71

Total	40,9°T	11,525	752	Sh0.0	88.0	7 <u>.</u> . 7
paiuL	505	698°T	LS	0.042	85.0	87.9
puopag	6E t7	866.5	607	980°0	62.0	E8 ' 9
degi ^H	996	891.7	Т9\$	090°0	78.0	7.42
gnilisM	Number	IstoT aysq blaila	TetoT ts9vasH	Average Daily Bag	Average Season Bag	Average Days Afield

Summary Deer Bow and Gun (Hunters not additive) Harvest Mail Survey $1970-71\,$ Table 9.

Based on 3,205 Usable Responses

Mailing	Number Hunters	Total Days	Total Harvest	Average Daily Bag	Average Season Bag	Average Days Afield
First	972	7,807	376	0.048	0.39	8.03
Second	7 7 7	3,150	113	980.0	0.25	7.09
Third	203	1,449	09	0.041	0.30	7.24
Total	1,619	12,406	5 49	110.0	0.34	7.66

Summary Turkey Harvest Mail Survey 1970-71 Based on 3,205 Usable Responses Table 10.

Mailing	Number Hunters	Total Days Afield	Total Harvest	Average Daily Bag	Average Season Bag	Average Days Afield	
				-			
First	202	6 44	29	0.071	0.33	19.4	
Second	8 2	358	50	0.065	0.24	4.21	
Third	88	153		0.027	0.25	5.46	
Total	315	1,455	16	0.065	0.30	4.62	

Summary Squirrel Harvest Mail Survey 1970-71 Based on 3,205 Usable Responses

			the state of the s		And the second s	
Mailing	Number Hunters	Total Days Afield	Total Harvest	Average Daily Bag	Average Season Bag	Average Days Afield
First	1,166	8,766	17,903	2.04	15.35	7.52
Second	595	3,979	8,312	2.09	13.97	
Third	237	2,063	4,129	2.00	17.42	8.70
Total	1,998	14,808	30,344	2.05	15.19	7.47

Table 12. Summary Duck Harvest Mail Survey 1970-71

Based on 3,205 Usable Responses

Number Hunters	Total Days Afield	Total Harvest	Average Daily Bag	Average Season Bag	Average Days Afield
319	1,500	2,414	1.61	7.57	4.70
110	525	894	1.70	8.13	4.77
51	258	341	1.32	6.70	5.06
480	2,283	3,649	1.60	7.60	4.76
	319 110 51	Number Days Hunters Afield 319 1,500 110 525 51 258	Number Hunters Days Afield Total Harvest 319 1,500 2,414 110 525 894 51 258 341	Number Hunters Days Afield Total Harvest Daily Bag 319 1,500 2,414 1.61 110 525 894 1.70 51 258 341 1.32	Number Hunters Days Afield Total Harvest Daily Bag Season Bag 319 1,500 2,414 1.61 7.57 110 525 894 1.70 8.13 51 258 341 1.32 6.70

Table 13. Summary Woodcock Harvest Mail Survey 1970-71.

Based on 3,205 Usable Responses

70.4	72.7	67.I	727	2.04	OOT	Total
09.2	. 00°01	67.I	09	82	S	baidT
85.8	TE • 6	27.5	242	88	56	Second
4.22	08.3	6 th • T	<u>5</u> 87	162	69	taríЯ
Average Days Afield	Average Bag	Average Daily Bag	LstoT tsevasH	LstoT aysU bleila	Number sastnuH	gnilisM

Table 14. County Summary of Hunting Pressure

Based on 2,633 Usable Responses

County	Total Man Days Hunted	Total Hunter Response	Percent Hunting Pressure
Adams Alcorn Amite Attala Benton Bolivar Calhoun Carroll Chickasaw Choctaw Claiborne Clarke Clay Coahoma Copiah Covington DeSoto Forrest Franklin George Greene Grenada Hancock Harrison Hinds Holmes Humphreys Issaquena Itawamba Jackson Jasper Jefferson Jefferson Jones	883 26994 51994 51994 51994 51994 51994 61944 61	58 208 46 46 47 40 40 40 40 40 40 40 40 40 40 40 40 40	1.63% 1.29% 1.07% 1.07% 1.07% 1.025% 1.04% 1.094% 1.094% 1.094% 1.287% 1.299% 1.287% 1.14% 1.287% 1.14
Kemper Lafayette Lamar Lauderdale Lawrence Leake Lee Leflore	1,033 1,068 477 1,016 270 560 770	110 52 80 35 42 72 71	1.97% .88% 1.87% .49% 1.03% 1.42%

Table 14. County Summary of Hunting Pressure (Continued)

Based on 2,633 Usable Responses

County .	Total Man Days Hunted	Total Hunter Response	Percent Hunting Pressure
Lincoln Lowndes Madison Marion Marshall Monroe Montgomery Neshoba Newton Noxubee Oktibbeha Panola Pearl River Perry Pike Pontotoc Prentiss Quitman Rankin Scott Sharkey Simpson Smith Stone Sunflower Tallahatchie Tate Tippah Tishomingo Tunica Union Walthall Warren Washington Wayne Webster Wilkinson Winston Yalobusha Yazoo	532 753 624 930 930 932 739 932 739 622 834 94 837 632 834 937 530 837 539 632 632 632 632 632 632 632 632 632 632	40 428 7537 75458 4654 71156 439 439 436 437 437 437 437 437 437 437 437 437 437	98% 1.39% 1.73% 1.83% 1.883% 1.883% 1.888% 1.189% 1.199% 1.29% 1.29% 1.29% 1.31% 1.29% 1.31% 1.29% 1.31% 1.31% 1.31% 1.31% 1.31% 1.31% 1.31% 1.31% 1.31% 1.31% 1.31% 1.31% 1.32% 1.33% 1.38% 1.3

Table 15. District Summary of **Hunting** Pressure

Based on 2,633 Usable Responses

	*,		
Game Ranagement District	Total Man Days Hunted	Total Hunter Hesponse	Percent Hunting Pressure
District I	8,468	734	15.64%
District II	8,783	773	16.22%
District III	7,948	709	14.68%
District IV	9,581	901	17.70%
District V	8,663	769	16.00%
District VI	10,681	974	19.76%
Total	54,124	4,860	100.00%
		•	

Table 16. Licensees Who Have Hunted or Trapped Other Species in Mississippi

Usable Responses: 3,205

Species	Respondents Who Hunted or Trapped	Percent	Respondents Who Did Not Per Hunt or Trap	cent
		and the second s		
Fox	270	8.42%	2,935 91	. 56%
Crow	260	8.11%	2,945 91	. 89%
Raccoon	494	15.28%	2,711 84	.56%
Opossum	240	7.50%	2,965 92	2.51%
Beaver	73	2.28%	3,132 97	.72%
Bobcat	146	4.41%	3,059 95	. 44%
Geese	39	1.22%	3,166 98	.78%
Other	45	1.39%	3,160 98	.59%

Table 17. Licensees Who Have Hunted or Trapped Other Species in Mississippi

Expanded Summary

Total Licensees: 240,479

Species	Estimated Hunters	Standard	Standard Error	
7 7 P		garagen agreement and a second a		
Fox	19,907	1,574	7.9%	
Crow	18,326	1,561	8.5%	
Raccoon	36,737	2,321	6.3%	
Opossum	17,820	1,909	0.7%	
Beaver	4,909	644	3.1%	
Bobcat	10,790	882	8.2%	
Geese	2,692	480	7.8%	
Other	3,090	472	5.3%	

Table 18. Confidence Limits at 95% Probability Level

Expanded Estimate of Licensees Who Have Hunted Or Trapped Other Species in Mississippi (Variation From the Mean)

Species	Estimated Hunters
Fox	19,907 ± 3,085
Crow	18,326 ± 3,059
Raccoon	36,737 ± 4,549
Opossum	17,820 ± 3,741
Beaver	4,909 ± 1,262
Bobcat	10,790 ± 1,728
Reese	2,692 ± 940
other	3,090 ± 925

Table 19. Expanded Summary 1970-71 Mississippi Post-Season Game Harvest Mail Survey

Total Licensees: 240,479

Species	Total Hunters	Percent of Total Permittees	Total Days Afield	Total Harvest	Average Daily Bag'	Average Season Bag	Average Days Afield
Mourning Dove	81,904	34.06	357,394	2,133,530	5.97	26.05	4.36
Quail	75,411	31.36	639,427	2,173,500	3.40	28.82	8.48
Rabbit	91,442	38.02	702,847	1,059,120	1.51	11.58	7.69
Deer - Archery Season	10,296	4.28	62,543	1,608	0.026	0.16	6.07
Deer - Gun Season	118,935	49.46	853,984	35,886	0.042	0.30	7.18
Deer - Bow or Gun (Hunters not additive)	119,939	49.88	911,803	37,353	0.041	0.31	7.60
Turkey	22,548	9.38	103,165	6,355	0.062	0.28	4.58
Squirrel	151,046	62.81	1,134,180	2,298,120	2.03	15.21	7.51
Duck	33,217	13.81	159,080	254,236	1.60	7.65	4.79
Woodcock	6,844	2.85	26,839	54,862	2.04	8.02	3.92

Table 20. Mississippi Post-Season Game Harvest Mail Survey 1970-71

Expanded Estimate of Total Hunters

Species Total	Number of	Hunters	Standard	Error
Mourning Dove	81,904		3,697	4.5%
Quail	75,411		2,619	3.5%
Rabbit	91,442		3,070	3.4%
Deer - Archery Season	10,296		883	8.6%
Deer - Gun Season	118,935		2,412	2.0%
Deer by Either Bow or Gun (Hunters not additive)	119,939		2,494	2.1%
Turkey	22,548		839	3.7%
Squirrel	151,046		3,678	2.4%
Duck	33,217	•	2,422	7.3%
Woodcock	6,844		932	13.6%

Table 21. Mississippi Post-Season Game Harvest Mail Survey 1970-71

Expanded	Estimate	of	Total	Days	Afield
----------	----------	----	-------	------	--------

Species	Total Days Afield	Standard Error
Mourning Dove	357,394	18,816 5.3%
Quail	639,427	41,119 6.4%
Rabbit	702,847	39,920 5.7%
Deer - Archery Season	62,543	2,671 4.3%
Deer - Gun Season	853,984	13,898 1.6%
Deer by Either Bow or Gun (Hunters not additive	911,803	13,601 1.5%
Turkey	103,165	7,683 7.4%
Squirrel	1,134,180	34,455 3.0%
Duck	159,080	17,560 11.0%
Woodcock	26,839	4,400 16.4%

Table 22. Mississippi Post-Season Game Harvest Mail Survey 1970-71

Expanded Estimate of Total Harvest

Species	Total Harvest	Standard	Error
Mourning Dove	2,133,530	155,345	7.3%
Quail	2,173,500	124,347	5.7%
Rabbit	1,059,120	46,341	4.4%
Deer - Archery Season	1,608	37.2	2.3%
Deer - Gun Season	35,886	2,318	6.5%
Deer by Either Bow or Gun (Hunters not additive)	37,353	1,895	5.1%
Turkey	6,355	841	13.2%
Squirrel	2,298,120	33,697	1.5%
Duck	254,236	32,014	12.6%
Woodcock	54,862	8,333	15.2%

Table 23. Confidence Limits at 95% Probability Level

Expanded Estimate of Total Hunters, Total Days Afield, Total Harvest
(Variation from the Mean)

Species	Number of Hunters	Total Days Afield	Total Harvest
Mourning Dove	81,904 ± 7,246	357,394 ± 36,879	2,213,530 ± 304,476
Quail	75,411 ± 5,133	639,427 ± 80,593	2,173,500 ± 243,720
Rabbit	91,442 ± 6,017	702,804 ± 78,243	1,059,120 ± 90,828
Deer (Archery Season)	10,296 ± 1,731	62,543 ± 5,235	1,608 ± 729
Deer (Gun Season)	118,935 ± 4,727	853,984 ± 27,240	35,886 ± 4,543
Deer by Either Bow or Gun (Hunters not additive)	119,939 ± 4,888	911,803 ± 26,658	37,353 ± 3,714
Turkey	22,548 ± 1,644	103,165 ± 15,059	6,355 ± 1,648
Squirrel	151,046 ± 7,209	1,134,180 ± 67,532	2,298,120 ± 66,046
Duck	33,217 ± 4,747	159,080 ± 34,418	254,236 ± 62,747
Woodcock	6,844 ± 1,827	26,839 ± 8,624	54,862 ± 16,333

LITERATURE CITED

- Atwood, E. L. 1956. Validity of mail survey data on bagged water-fowl. J. Wildl. Mgmt., 20 (1): 1-16.
- Barick, Frank B., and T. Stuart Critcher. 1955. Statistical game kill surveys—some observations on five years of operation. 9th Ann. Meeting, S. E. Assn. Game and Fish Comm's. 43-47.
- Chapman, D. G., W. S. Overton, and A. L. Finkner. 1959. Methods of estimating dove kill. Mimeo Series No. 264, Institute of Statistics, North Carolina State College, Raleigh. 48 pp.
- Cronan, J. M. 1960. Prestige and memory bias in hunter kill surveys. Paper presented at the Northeast Wildl Conf., Providence, R. I., January 1960.
- Durell, James S. 1967. A brief study of hunters and the owners of the land on which they hunt. Proc. 21st Ann. Conf. S. E. Assn. Game and Fish Comm's. 81-87.
- Eberhardt, Lee. 1961. Estimating the kill of game animals by licensed hunters. American Statistical Association Journal. June 1961: 396-397.
- Hayne, D. W. 1964. Investigation of mail survey reporting by water-fowl hunters. Report prepared for U. S. Fish and Wildlife Service, Bureau of Sport Fisheries and Wildlife, Patuxent Wildlife Research Center. Institute of Statistics, N. C. S., Raleigh. 24 pp.
- Hiltunen, Eero. 1952. On the reliability of the hunting license statistics. Suomen Riista. 7: 72-81.
- Kelly, William F. 1967. Statewide game harvest 1966-67. P-R Proj. W-34-15, Work Plan No. 2, Job No. 2-B. Alabama Dept. Conservation. Unpubl.
 - P-R Proj. W-34-16, Work Plan No. 2, Job No. 2-B.
 Alabama Dept. conservation. Unpubl.
- Leopold, Aldo. 1929. Report on a game survey of Mississippi. Submitted to the Game Restoration Committee, Sporting Arms and Ammunition Manufacturer's Institute. February 1, 1929.